


Specimen Number	Specimen Type <b>Peripheral Blood</b>	Control Number	Account Number	Account Phone Number	Route	
Patient Last Name						
Patient First Name <b>Raven</b>	Patient Middle Name					
Patient SS#	Patient Phone					Total Volume
Age (Y/M/D) <b>25 y.o.</b>	Date of Birth					Sex <b>Female</b>
Patient Address		<b>Indication: Hemophilia B carrier</b> <b>Family History: Family history of uncontrolled bleeding</b> <b>Ethnicity: Native American, Navajo Tribe</b>				
Date and Time Collected	Date Entered	Date and Time Reported	Physician Name <b>Jane Ferreiro, MD</b>	NPI	Physician ID	

<b>Hemophilia Mutation Evaluation</b>	Tests Ordered
General Comments <b>Please send a copy of the final report to the Molecular Science/M1 Training office via Fax at (202) 555-1212</b>	

## Clinical test results for DNA Hemophilia Mutation Evaluation

GENE	TEST RESULTS	EXPLANATION
<b>F8 (Xq28)</b>	Arg15 Arg15Ter Heterozygous Carrier Status	<p><b>This result confirms the status of carrier for Hemophilia A.</b> This result should be interpreted in the context of clinical presentation and results of other laboratory tests (e.g., APTT, Factor VIII Activity, etc.).</p> <p>A PCR/sequencing study has detected one copy of the Arg15Ter (F8:g.5214C&gt;T, c.43C&gt;T or p.Arg15Ter) variation. The Arg15Ter variation is a C to T change at nucleotide position 5214 of the F8 gene and 43 of the F8 mRNA transcript. This forms a premature stop (termination) codon at amino acid position 15 resulting in an abnormally short or truncated protein.</p> <p>Female carriers of Hemophilia A have one normal X chromosome and one abnormal X chromosome. The normal X chromosome produces a certain amount of factor IX clotting factor. This protects carriers from the most severe form of hemophilia, however the levels of circulating clotting factors among carriers are very wide.</p>
<b>F9 (Xq27.1)</b>	Negative	

### INDICATIONS FOR TESTING

Individuals with a diagnosis of hemophilia B, appropriate at-risk female relatives of probands with identified mutations, and hemophilia B carriers with genetic counseling, are candidates for testing.

### METHODOLOGY

Factor IX sequencing: All coding exons (1-8) and associated intron junctions of the Factor IX gene are analyzed by direct DNA sequence analysis using an automated fluorescent sequencing machine. When a mutation is detected, confirmation is carried out on an independent amplification of PCR using a second prep (B-prep) by sequencing in the opposite direction. If no mutation is found, sequence analysis is performed in both directions.

### PERFORMANCE

Factor IX sequencing: From previous experience, we have been able to detect factor IX gene mutations in about 99% of individuals with the diagnosis of hemophilia B with specificity of mutation detection in probands and carrier detection is also estimated to be greater than 99%.

## LIMITATIONS

The sequence analysis will not detect mutations located in regions of the Factor IX gene that are not analyzed (non-coding exon regions, intron regions other than the splice junctions, and upstream and downstream regions). The sequencing method also will not detect gross genetic alterations including most duplications, inversions, or deletions (in females). Some sequence alterations that may be detected (such as those causing missense or synonymous changes) will be of unknown clinical significance.

## CLINICAL DESCRIPTION

Hemophilia B is characterized by deficiency in factor IX clotting activity that results in prolonged oozing after injuries, tooth extractions, or surgery, and delayed or recurrent bleeding prior to complete wound healing. This is an X-linked recessive bleeding disorder with an incidence of about 1 per 30,000 live male births. Hemophilia B affects males, however, all male offspring will be normal, and although all female offspring will be obligatory carriers, they rarely have symptomatic bleeding. In contrast, female offspring of carriers of hemophilia B have a 50% chance of being carriers themselves, and each male offspring has a 50% chance of being affected.

The age of diagnosis and frequency of bleeding episodes are related to the level of factor IX clotting activity. In severe hemophilia B, spontaneous joint or deep-muscle bleeding is the most frequent symptom. Individuals with severe hemophilia B are usually diagnosed during the first two years of life; without prophylactic treatment, they may average up to two to five spontaneous bleeding episodes each month. Individuals with moderate hemophilia B seldom have spontaneous bleeding; however, they do have prolonged or delayed oozing after relatively minor trauma and are usually diagnosed before age five to six years; the frequency of bleeding episodes varies from once a month to once a year. Individuals with mild hemophilia B do not have spontaneous bleeding episodes; however, without pre- and post-operative treatment, abnormal bleeding occurs with surgery or tooth extractions; the frequency of bleeding may vary from once a year to once every ten years. Individuals with mild hemophilia B are often not diagnosed until later in life. In any individual with hemophilia B, bleeding episodes may be more frequent in childhood and adolescence than in adulthood. Approximately 10% of carrier females are at risk for bleeding (even if the affected family member has mild hemophilia B) and are thus symptomatic carriers, although symptoms are usually mild. After major trauma or invasive procedures, prolonged or excessive bleeding usually occurs, regardless of severity.

## REFERENCES

4. Yoshitake S, Schach BG, Foster DC, et al: Nucleotide sequence of the gene for human factor IX (antihemophilic factor B). *Biochemistry* 1985 July 2;24(14):3736-3750
5. Giannelli F, Green PM, Sommer SS, et al: Haemophilia B: database of point mutations and short additions and deletions-eighth edition. *Nucleic Acids Res* 1998 Jan 1;26(1):265-268
6. Ketterling RP, Bottema CD, Phillips JA 3rd, et al: Evidence that descendants of three founders constitute about 25% of hemophilia B in the United States. *Genomics* 1991 Aug;10(4):1093-1096